

Paradoxical Associations Between Familial Affective Responsiveness, Stress, and Amygdala Reactivity

Madeline J. Farber, Adrienne L. Romer,
M. Justin Kim, and Annchen R. Knodt
Duke University

Nourhan M. Elsayed
Duke University School of Medicine

Douglas E. Williamson
Duke University School of Medicine and Durham VA Medical
Center, Durham, North Carolina

Ahmad R. Hariri
Duke University

Studies of early life extremes such as trauma, abuse, and neglect highlight the critical importance of quality caregiving in the development of brain circuits supporting emotional behavior and mental health. The impact of normative variability in caregiving on such biobehavioral processes, however, is poorly understood. Here, we provide initial evidence that even subtle variability in normative caregiving maps onto individual differences in threat-related brain function and, potentially, associated psychopathology in adolescence. Specifically, we report that greater familial affective responsiveness is associated with heightened amygdala reactivity to interpersonal threat, particularly in adolescents having experienced relatively low recent stress. These findings extend the literature on the effects of caregiving extremes on brain function to subtle, normative variability but suggest that presumably protective factors may be associated with increased risk-related amygdala reactivity. We consider these paradoxical associations with regard to studies of basic associative threat learning and further consider their relevance for understanding potential effects of caregiving on psychological development.

Keywords: caregiving, fMRI, amygdala, threat processing, stress

Supplemental materials: <http://dx.doi.org/10.1037/emo0000467.supp>

A wealth of empirical research has demonstrated that the quality of caregiving predicts later psychosocial outcomes, in part, by

shaping the development of brain circuits supporting emotional behavior (Belsky & de Haan, 2011; Callaghan & Tottenham, 2016; Cicchetti & Curtis, 2015). However, much of this research has focused on caregiving extremes, such as trauma, abuse, and neglect (Gee et al., 2013; Herringa et al., 2013; Malter Cohen et al., 2013; McLaughlin, Peverill, Gold, Alves, & Sheridan, 2015; Pechtel, Lyons-Ruth, Anderson, & Teicher, 2014; Sheridan, Fox, Zeanah, McLaughlin, & Nelson, 2012; Tottenham, 2012; Tottenham et al., 2010). Such a focus ignores the potential impact of normative variability in caregiving on neurodevelopment and behavior. A rich literature has demonstrated lasting positive effects of normative caregiving including secure attachment and positive parenting (i.e., high care, low control) on behavioral development (Bowlby, 1978; Bowlby & Zeanah, 1988; Canetti, Bachar, Galili-Weisstub, De-Nour, & Shalev, 1997; Francis & Meaney, 1999; Overbeek, ten Have, Vollebergh, & de Graaf, 2007). Despite this literature, comparatively little is known about the potential impact of normative variability in caregiving on the underlying brain circuitry implicated in both normal and abnormal behavior (Rohmund et al., 2016; Tan et al., 2014; Whittle et al., 2009).

Here, we seek to address this research gap by examining how normative variability in caregiving within the family affects individual differences in threat-related amygdala reactivity in a large cohort of adolescents ages 12–15 years. We focus on threat-related

This article was published Online First July 12, 2018.

Madeline J. Farber, Adrienne L. Romer, M. Justin Kim, and Annchen R. Knodt, Laboratory of Neurogenetics, Department of Psychology & Neuroscience, Duke University; Nourhan M. Elsayed, Translational Center for Stress-Related Disorders, Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine; Douglas E. Williamson, Translational Center for Stress-Related Disorders, Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine, and Durham VA Medical Center, Durham, North Carolina; Ahmad R. Hariri, Laboratory of Neurogenetics, Department of Psychology & Neuroscience, Duke University.

The main funding for this study was through Duke University and the National Institute on Alcohol Abuse and Alcoholism Grant R01AA016274 (Douglas E. Williamson). The Laboratory of Neurogenetics received support from Duke University as well as National Institutes of Health Grants R01DA033369, R01DA03157, and R01AG049789. We thank the Teen Alcohol Outcomes Study participants as well as the staff of the Laboratory of Neurogenetics and the Translational Center for Stress-Related Disorders.

Correspondence concerning this article should be addressed to Madeline J. Farber, Laboratory of Neurogenetics, Department of Psychology & Neuroscience, Duke University, Box 104151, Durham, NC 27708. E-mail: madeline.farber@duke.edu

amygdala reactivity because of converging evidence that this specific aspect of brain function not only is affected by extremes of caregiving (Gee et al., 2013; Herringa et al., 2013; Malter Cohen et al., 2013; McLaughlin et al., 2015; Pechtel et al., 2014; Sheridan et al., 2012; Tottenham, 2012; Tottenham et al., 2010) but also represents a biomarker of future risk for stress-related depression and anxiety (Admon et al., 2013; Etkin & Wager, 2007; Schlund & Cataldo, 2010; Swartz, Knodt, Radtke, & Hariri, 2015). Given the body of work highlighting the potential of stressful life events to trigger the development of anxiety and depression in part through amygdala function (Kessler et al., 2005; Paus, Keshavan, & Giedd, 2008; Swartz, Williamson, & Hariri, 2015), we additionally investigate whether stress plays a moderating role in the relationship between caregiving and amygdala reactivity to threat.

In contrast to the extant literature, which represents a nearly exclusive focus on the effects of maternal caregiving (Caldji, Diorio, & Meaney, 2003; Francis & Meaney, 1999; Gee et al., 2013; Romund et al., 2016; Tan et al., 2014; Tottenham, Shapiro, Telzer, & Humphreys, 2012; Whittle et al., 2009), we examine holistic family functioning to help inform more general effects of the caregiving environment on behaviorally and clinically relevant brain function. To this end, given the complex interpersonal dynamics at play within the family system, we focus on amygdala reactivity to interpersonal threat cues as specifically indexed by angry facial expressions (Adams, Gordon, Baird, Ambady, & Kleck, 2003).

Our analyses centered on data available for a subset of participants having completed the Teen Alcohol Outcomes Study (TAOS; Birmaher et al., 2000), originally developed to examine the impact of familial loading for depression on the risk to develop alcohol use disorders. Study participants were characterized as either “high risk” (having a first- and second-degree relative with major depressive disorder [MDD]) or “low risk” (having no first-degree and minimal second-degree relatives [$<20\%$] with MDD; Williamson, Birmaher, Axelson, Ryan, & Dahl, 2004). The broader caregiving environment was indexed through the lens of family functioning as measured by the general functioning and affective responsiveness subscales of the Family Assessment Device (FAD; Epstein, Baldwin, & Bishop, 1983). Amygdala reactivity to explicit, interpersonal threat and to implicit, environmental threat as communicated by angry and fearful facial expressions, respectively, was measured using task-based functional magnetic resonance imaging (fMRI).

We hypothesized that lower scores on the FAD affective responsiveness subscale, which reflects appropriate expression and recognition of emotion through warmth, care, and affection, would be associated with decreased amygdala reactivity specifically to angry facial expressions as canonical examples of explicit interpersonal threat. Our hypothesis is based on the assumption that interpersonal conflict is less likely to be experienced when family members appropriately express emotions and that this positive modeling of emotional experiences would thus be reflected in lower amygdala reactivity to signals of such conflict. As an extension of this primary hypothesis as well as the literature reviewed earlier, we further explored the following related questions. Are significant associations with amygdala reactivity specific to affective responsiveness and not more general family functioning? Moreover, are significant associations independent of early life stress, contemporaneous symptoms of depression and

anxiety, and broad familial risk for depression? And, last, are significant associations moderated by recent stressful life events?

Method

Participants

Study participants were recruited into two groups: high risk, wherein a first- and second-degree relative has a history of MDD, or low risk, wherein no first-degree and minimal second-degree relatives ($<20\%$) have such a history. Low-risk participants further could not meet criteria for any psychiatric disorder or substance use disorder; however, a diagnosis of an anxiety disorder was permitted in the high-risk participants because early onset anxiety is frequently a precursor to depression (Goodwin, Ferguson, & Horwood, 2004; McLaughlin & King, 2015). Mental health in all participants was assessed through structured clinical interviews with the adolescent and parent separately using the Kiddie Schedule for Affective Disorders and Schizophrenia for School-Age Children—Present and Lifetime version (Kaufman et al., 1997). All procedures were approved by the Institutional Review Board at the University of Texas Health Sciences Center San Antonio, and participants and their parent provided written informed assent–consent before participation. Additional recruitment details have been described in detail elsewhere (Bogdan, Williamson, & Hariri, 2012; Swartz et al., 2017; White et al., 2012).

Of the 331 participants completing the parent study (the Teen Alcohol Outcomes Study), data for our present analyses were available for 232 participants (120 high risk, 112 low risk). Of these 331 participants, data were excluded for 29 due to problems with the scan or raw data (e.g., ending the scan early, poor coverage of the amygdala, gross anatomical abnormalities, scanner artifacts). Data from another seven high-risk participants were excluded because they had a diagnosis of MDD before the scan and from another five low-risk participants because they had an anxiety disorder before the scan. Functional MRI data from the remaining 290 participants underwent preprocessing, and subsequently, another 53 participants were excluded based on quality control criteria for the processed data (see the Functional MRI section for further details). Last, data from another five participants were excluded for poor accuracy on the fMRI task. The final analysis sample of 232 participants was 50.4% female and approximately 58% Caucasian; 26% Hispanic; 5% African American; and 12% Indian, Asian, Pacific Islander, or multiple races. There was no difference in sex ratio, $\chi^2(1, N = 232) = .018, p = .892$, or the distribution of race–ethnicity, $\chi^2(4, N = 232) = 2.413, p = .660$, between the high-risk and low-risk groups. There were no differences between included and excluded participants in age, gender, family functioning, childhood trauma, depressive symptoms, anxiety symptoms, or stressful life events.

Behavioral Measures

Family Assessment Device (FAD). Family functioning was assessed using the FAD, a 60-item scale completed by both child and parent, which yields seven scales of family functioning: six subscales (affective responsiveness, problem solving, communication, roles, affective involvement, behavior control) and one global

measure (general functioning). Family members rate how each item describes their family (no explicit time window provided) by selecting 1 (*strongly agree*), 2 (*agree*), 3 (*disagree*), or 4 (*strongly disagree*), 1 reflecting healthy functioning and 4 reflecting unhealthy functioning (Epstein, Baldwin, & Bishop, 1983; Stein et al., 2000). The use of FAD subscales is normed and well-validated in the literature. The FAD has been used in a variety of research contexts and has shown good reliability for all subscales (Cronbach's alphas: affective responsiveness = .73, general functioning = .83, problem solving = .74, communication = .73, roles = .57, affective involvement = .76, behavior control = .70; Epstein et al., 1983; Kabacoff, Miller, Bishop, Epstein, & Keitner, 1990). Although there is no total score for the FAD, the general functioning subscale is often considered a global score in interpretation.

Given the utility of these subscales as independent measures, we focused on the affective responsiveness subscale because items from this subscale best represent warmth, care, and affection provided by caregivers within the broad family dynamic. Examples include "We are reluctant to show our affection for each other" and "We express tenderness." We addressed the possible unique associations of this caregiving feature with amygdala reactivity by also examining post hoc correlations with the *general functioning* subscale, which measures the overall level of family functioning with items such as "In times of crisis we can turn to each other for support" and "We don't get along well together," as well as the five remaining FAD subscales. In all analyses, we utilized child- and not parent-reported FAD scores because we expected that children's perception of their familial environment would be more impactful than would the parents' perception, and we wanted to better approximate scores from the Parental Bonding Instrument, which is a more commonly used inventory of normative caregiving that is child-reported but unavailable in the present study.

Mood and Feelings Questionnaire (MFQ). Depressive symptoms were assessed using the MFQ, a widely used measure in children and adolescents (Messer et al., 1995). The MFQ assesses depressive symptoms in the last 2 weeks. We used the child-reported short (13-item) summary score in our analyses.

Screen for Child Anxiety Related Disorders (SCARED). Anxiety symptoms were assessed using the SCARED, a widely used measure of anxiety symptoms in children and adolescents (Birmaher et al., 1999). The SCARED assesses anxiety symptoms across the previous 3 months. We use the child-reported SCARED in our analyses. The MFQ and SCARED have both shown good reliability across studies (.90 and .60–.90, respectively; Angold et al., 1995; Birmaher et al., 1999).

Childhood Trauma Questionnaire (CTQ). Early life stress was assessed using the CTQ, a widely used measure of childhood trauma and early life adversity (Bernstein et al., 2003). The CTQ assesses trauma in childhood broadly without an explicit time window. This measure has shown good reliability in normative community samples (.79–.86; Scher, Stein, Asmundson, McCreary, & Forde, 2001) and in psychiatric adolescent samples (.96–.97; Bernstein, Ahluvalia, Pogge, & Handelsman, 1997).

Stressful Life Events Schedule (SLES). Recent stress was measured using the SLES, a structured interview designed to assess for life stressors relevant to children and adolescents (Williamson et al., 2003). The SLES assesses stressful life events in the

past year. Although the SLES does peripherally query some life stressors pertaining to family members, this is not its primary focus, nor does it thoroughly index family environment or parenting style; it can therefore be considered a measure of external stressors independent of those indexed via the FAD. The SLES has shown good interrater ($\kappa = .67$) and test–retest ($\kappa = .68$) reliability (Williamson et al., 2003).

Functional MRI

Participants performed an emotional face matching task that has been shown to consistently elicit robust amygdala activity in numerous studies of adults and adolescents (Bogdan et al., 2012; Swartz et al., 2017; White et al., 2012). Task blocks consisted of matching angry facial expressions and fearful facial expressions, whereas control blocks consisted of matching geometric shapes (Bogdan et al., 2012; Swartz, Williamson, et al., 2015; Swartz et al., 2017; White et al., 2012). Further task details are reported in the [online supplemental materials](#).

Analyses were conducted using the general linear model of SPM8 (Penny, Friston, Ashburner, Kiebel, & Nichols, 2011). Following preprocessing, linear contrasts employing canonical hemodynamic response functions were used to estimate main effects of expression for each individual. Individual contrast images were then entered in second-level random effects models to determine mean condition-specific regional responses using one-sample *t* tests. We extracted parameter estimates from functional clusters within anatomically defined amygdala regions of interest (Automated Anatomical Labeling atlas) at $p < .05$ family-wise error (FWE) corrected across the search volumes for the contrast of angry blocks > control blocks and fearful blocks > control blocks (Swartz, Williamson, et al., 2015). Further quality assurance and extraction details are reported in the [online supplemental materials](#).

Statistical Analyses

Mean individual contrast-related blood oxygen level-dependent (BOLD) activation values from functional clusters were entered into second-level analyses in SPSS, Version 24. In all analyses, we controlled for participant age, sex, and risk group. Additionally, we controlled for participant CTQ scores in all analyses because the relationship between childhood trauma and amygdala reactivity is well documented (Gee et al., 2013; Heringa et al., 2013; Malter Cohen et al., 2013; McLaughlin et al., 2015; Pechtel et al., 2014; Sheridan et al., 2012; Tottenham, 2012; Tottenham et al., 2010) and we aimed to probe the unique contribution of specific dimensions of family functioning.

First, to test our primary hypothesis, we ran a linear regression of the FAD affective responsiveness subscale with extracted BOLD values for the contrast of angry expressions greater than shapes. Second, to confirm that the significant association between familial affective responsiveness and amygdala reactivity was not driven by concurrent depressive and anxiety symptoms, we duplicated regression analyses from Step 1, this time controlling for MFQ and SCARED scores in addition to our other covariates. Last, we conducted linear regression analyses to investigate potential direct effects of recent stress (SLES) and interactions of SLES with FAD on amygdala reactivity to angry facial expres-

sions. We then conducted post hoc partial correlation analyses within the high- and low-stress groups to test for significance of the simple slopes.

To probe the specificity of the association of our primary hypothesis (i.e., FAD affective responsiveness and amygdala reactivity to angry facial expressions), we conducted two sets of post hoc analyses: (a) linear regressions analyses of the FAD affective responsiveness subscale with extracted BOLD values for amygdala reactivity to fearful expressions and (b) linear regression analyses of the FAD general functioning scale and remaining FAD subscales (problem solving, communication, roles, affective involvement, and behavior control) with extracted BOLD values for amygdala reactivity to angry expressions.

Results

Participant Characteristics

Demographic and behavioral characteristics of our analysis sample are detailed in Table 1. High- and low-risk groups significantly differed in symptoms of depression, $t(200.4) = 2.405, p = .019$, and in early life stress, $t(212.2) = 2.228, p = .027$. However, the high- and low-risk groups did not differ in FAD affective responsiveness, $t(229) = -.382, p = .703$. Groups also did not differ in anxiety symptoms, $t(217.3) = 1.148, p = .252$, or recent life stress, $t(210.3) = 1.234, p = .219$. There were no significant differences between groups on possible neuroimaging confounds including task performance as measured by accuracy, $t(225) = -.230, p = .818$, and response time, $t(219) = -.552, p = .582$, or in task-related movement as indexed by head displacement, $t(230) = -.609, p = .543$. Further details regarding FAD subscale distribution and between-subscales correlations within our sample are reported in the [online supplemental materials](#). Zero-order correlations between our primary behavioral measures of interest (FAD, CTQ, SLES, MFQ, SCARED) are reported in Table 2.

Family Functioning and Amygdala Reactivity

Across all participants, there was robust bilateral amygdala reactivity to both angry and fearful facial expressions (see Figure

1). Linear regression analyses using extracted BOLD parameter estimates from clusters exhibiting main effects of expression revealed a significant association between FAD affective responsiveness and amygdala reactivity to angry facial expressions. Contrary to our hypothesis, participants who reported better affective responsiveness (i.e., lower scores) exhibited significantly higher reactivity to angry facial expressions in the left ($\beta = -.190, p = .015$) but not right ($\beta = -.076, p = .328$) hemisphere (see Figure 2). This association was independent of broader familial risk for depression, because there was no statistically significant interaction between affective responsiveness and risk-group status on reactivity ($\beta = .026, p = .906$).

Subsequent analyses revealed that the association between affective responsiveness and amygdala reactivity to angry facial expressions was further independent of contemporaneous symptoms of anxiety and depression ($\beta = -.187, p = .017$). Neither depression ($\beta = -.088, p = .341$) nor anxiety ($\beta = -.084, p = .308$) significantly contributed to amygdala reactivity to angry facial expressions. There were no significant main effects of FAD affective responsiveness on symptoms of depression ($\beta = -.006, p = .932$) or anxiety ($\beta = .084, p = .277$). Likewise, there was no significant interaction effect of left amygdala reactivity to angry facial expressions by FAD affective responsiveness on symptoms of depression ($\beta = .026, p = .671$) or anxiety ($\beta = .007, p = .921$).

Last, regression analyses revealed a significant interaction of recent life stress and FAD affective responsiveness ($\beta = .142, p = .050$), such that the association between affective responsiveness and amygdala reactivity to angry facial expressions was moderated by the experience of recent life stress (see Figure 3). Specifically, the association between healthier affective responsiveness and higher left amygdala reactivity to angry facial expressions was significant for participants reporting lower ($r = -.244, p = .014$) but not higher ($r = -.052, p = .614$) recent life stress. There was no significant main effect of SLES on amygdala reactivity to angry facial expressions ($\beta = -.093, p = .186$). Details of independent regression coefficients yielded from this set of analyses are reported in the [online supplemental materials](#).

Table 1
Demographic and Behavioral Characteristics of Analysis Sample by Risk Group

Variable	High risk ($n = 120$)		Low risk ($n = 112$)		Group comparisons	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>t</i>	<i>p</i>
Age	13.61	1.01	13.56	.94	.416	.678
FAD aff resp	16.69	2.72	16.84	2.96	.483	.630
MFQ	4.06	4.20	2.96	2.60	2.405	.017*
SCARED	16.64	11.46	15.12	8.51	1.148	.252
CTQ	34.55	6.85	32.59	6.01	2.228	.027*
SLES	13.40	10.22	12.20	8.44	1.234	.219
Mean accuracy (%)	95.72	17.42	96.23	15.92	-.237	.818
Mean response time (ms)	1,288.61	240.85	1,306.66	245.35	-.522	.582
Mean head displacement (mm)	.04	.01	.04	.02	-.609	.543

Note. FAD aff resp = affective responsiveness subscale of the Family Assessment Device; MFQ = Mood and Feelings Questionnaire; SCARED = Screen for Child Anxiety Related Disorders; CTQ = Childhood Trauma Questionnaire; SLES = Stressful Life Events Schedule.

* $p < .05$.

Table 2
Pearson Correlations for Primary Behavioral Measures of Interest

Variable	1	2	3	4	5	6
1. Age	—	-.049	.027	-.010	-.055	.057
2. FAD aff resp	-.049	—	.201**	.111	.486**	.105
3. MFQ	.027	.201**	—	.534**	.456**	.398**
4. SCARED	-.010	.111	.534**	—	.140*	.261**
5. CTQ	-.055	.486**	.456**	.140*	—	.274**
6. SLES	.057	.105	.398**	.261**	.274**	—

Note. FAD aff resp = affective responsiveness subscale of the Family Assessment Device; MFQ = Mood and Feelings Questionnaire; SCARED = Screen for Child Anxiety Related Disorders; CTQ = Childhood Trauma Questionnaire; SLES = Stressful Life Events Schedule.

* $p < .05$. ** $p < .01$.

It is important to note that post hoc analyses revealed there were no significant associations between FAD affective responsiveness and amygdala reactivity to fearful facial expressions (left: $\beta = -.016$, $p = .838$; right $\beta = -.052$, $p = .495$). Moreover, post hoc analyses revealed there were no significant associations between amygdala reactivity to angry facial expressions and FAD general functioning or any of the five remaining FAD subscales (full statistics reported in the [online supplemental materials](#)).

Discussion

Our findings extend the literature on the impact of caregiving extremes on behaviorally relevant brain function by demonstrating a significant association between normative variability in family functioning and threat-related amygdala reactivity. Specifically, we found that even subtle differences in caregiving within the family map onto variability in amygdala reactivity to explicit signals of interpersonal threat (i.e., angry facial expressions). Contrary to our hypothesis, however, better familial affective responsiveness was associated with increased amygdala reactivity to interpersonal threat. This association was robust to the potential influence of participant sex, age, broad familial risk for depression, and early life stress, as well as contemporaneous symptoms of

depression and anxiety. Moreover, this association was moderated by the experience of recent stressful life events, wherein better affective responsiveness was associated with higher amygdala reactivity in participants reporting low but not high recent stress.

Our paradoxical associations are particularly interesting when conceptualizing family functioning as an interpersonal process. For example, in families in which emotions are not appropriately processed, children may frequently experience interpersonal conflict including anger between family members and thus may be desensitized to interpersonal threat cues, as evidenced by lower amygdala reactivity to angry facial expressions. Alternatively, it is possible that children raised by parents who more appropriately experience, express, and model their emotions are sheltered from interpersonal conflict, making them more sensitive to interpersonal threat cues and thus more likely to exhibit amygdala hyperreactivity to angry facial expressions. Consistent with this speculation, our findings revealed that the association between familial affective responsiveness and amygdala reactivity to angry expressions was specific to participants who reported having experienced relatively less recent life stress. Thus, adolescents who are raised in emotionally expressive, supportive family environments and




Figure 1. Threat-related amygdala reactivity. Mean reactivity to angry (Panel a) and fearful (Panel b) facial expressions within anatomically defined amygdala regions of interest ($p < .05$, family-wise error-corrected). Panel a: Left: cluster size = 169 voxels, peak voxel = $-22, -2, -18$; right: cluster size = 226 voxels, peak voxel = $20, -4, -16$. Panel b: Left: cluster size = 170 voxels, peak voxel = $-22, -2, -18$; right: cluster size = 224 voxels, peak voxel = $22, -4, -18$. Color (grayscale in the print edition) bar represents t scores. See the online article for the color version of this figure.




Figure 2. Family functioning and amygdala reactivity. Family Assessment Device (FAD) affective responsiveness scores are negatively correlated with left amygdala reactivity to interpersonal threat as indexed by angry facial expressions ($\beta = -.190, p = .015$). Of note, higher scores on the FAD reflect poorer functioning, whereas lower scores reflect better functioning.

who experience few external stressors may be more likely to exhibit neural hypersensitivity to interpersonal threat, perhaps reflecting their lack of exposure to conflict.

Of importance, there is support for our observed paradoxical findings from both basic biobehavioral and clinical developmental research. For example, work by Fanselow and Tighe (1988) demonstrated that the length of time between aversive shocks has a greater impact on freezing behavior in rats than does the frequency of shocks. Specifically, unpredictable, infrequent shocks result in stronger freezing behavior than do regular, frequent shocks. This pattern of potentiated aversive learning, which is mediated by amygdala circuitry, is more generally consistent with the increased effectiveness of spaced versus massed learning (Bouton, 2007). Thus, the observed amygdala hyperreactivity to interpersonal threat cues in adolescents from families with better affective responsiveness who experience lower recent stress may reflect heightened associative learning following less frequent, more unpredictable experiences of threat or conflict.

In parallel to this basic research, developmental clinical psychology studies have observed patterns between family functioning and threat-related biobehavioral processes similar to those we observed. For example, Andreotti and colleagues found that only young adults who reported lower levels of family conflict in early life exhibited increased cortisol reactivity to an acute laboratory stressor that subsequently predicted subliminal attentional bias to threat cues (Andreotti, Garrard, Venkatraman, & Compas, 2015).

Notably, the amygdala plays a significant role in both processing subliminal threat signals and triggering cortisol release via activation of the hypothalamic–pituitary–adrenal axis (Whalen & Phelps, 2009). More generally, parental overprotection, which may be characterized by either more restrictive and controlling (Parker, 1983) or indulgent (Green & Solnit, 1964) parenting has been associated with later psychopathology, including depression and anxiety disorders (Thomasgard & Metz, 1993). Consistent with these patterns, amygdala hyperreactivity is a core pathophysiological observation in these disorders (Etkin & Wager, 2007) and has emerged as a biomarker of future risk (Admon et al., 2009; Mattson, Hyde, Shaw, Forbes, & Monk, 2016; McLaughlin et al., 2014; Swartz, Knodt, et al., 2015).

It is interesting to further speculate that our observed associations between better familial affective responsiveness, less stress, and higher amygdala reactivity to angry facial expressions may actually be adaptive. Although we could not test this hypothesis in the present study, it is possible that we were observing the first half of a curvilinear, U-shaped relationship between familial affective responsiveness and amygdala reactivity. The TAOS data set consists of largely psychologically healthy adolescents. With the exception of a few low-risk participants with anxiety disorders, no participants in this sample had a clinical diagnosis at the time of scanning. Were we to duplicate the present analyses in a sample of matched adolescents with clinically diagnosed disorders, we might see an inverse relationship between familial affective responsive-




Figure 3. Family functioning, recent stress, and amygdala reactivity. The negative correlation between Family Assessment Device (FAD) affective responsiveness and amygdala reactivity to interpersonal threat was significant for only participants reporting relatively low levels of recent stressful life events as assessed by the Stressful Life Events Schedule ($\beta = .142, p = .050$; low stress: $r = -.244, p = .014$; high stress: $r = -.052, p = .614$). Of note, higher scores on the FAD reflect poorer functioning, whereas lower scores reflect better functioning.

ness and amygdala reactivity to interpersonal threat in which adolescents with psychiatric diagnoses reporting better familial affective responsiveness exhibit relatively decreased amygdala reactivity. Such speculation finds some support in the *attenuation hypothesis* (Susman, 2006), which states that at-risk samples may exhibit blunted emotion processing in some cases, particularly in considering the transmission of family aggression (Saxbe, Del Piero, Immordino-Yang, Kaplan, & Margolin, 2016). That said, we found no evidence for a differential association between amygdala reactivity and affective responsiveness as a function of risk-group status.

To our knowledge, this is one of few studies investigating the neural correlates of normative variability in caregiving in a non-clinical sample of adolescents. In addition to our current findings, Romund et al. (2016) have reported that increased maternal warmth and support are correlated with lower amygdala reactivity to fearful facial expressions among a healthy sample of adolescents, although they did not find an association with reactivity to angry facial expressions. The divergence of our associations with that of Romund et al.'s may reflect differences between broader, nonspecific familial dynamics and maternal-specific behaviors and explicit modeling of recent stress, as well as differences in the task-based amygdala reactivity elicited. Of note, unlike our conservative approach using unbiased amygdala reactivity values originating from our main effects of expression contrasts in our analyses of a much larger sample (232 vs. 83), Romund et al. did not find significant main effects of angry or fearful facial expressions on amygdala reactivity. Rather, they conducted more per-

missive correlation analyses identifying specific voxels within the amygdala that exhibited BOLD values correlated with their measures of maternal caregiving.

In another recent study, Thijssen et al. (2017) reported that insensitive parenting was associated with a more mature pattern of intrinsic connectivity between the amygdala and medial prefrontal cortex, which together function to integrate and regulate emotional experiences, in healthy children. And in an earlier study, Taylor, Eisenberger, Saxbe, Lehman, and Lieberman (2006) reported that lower family stress during childhood was associated with lower correlated activity between the amygdala and regulatory regions of the ventrolateral prefrontal cortex during affective labeling of angry and fearful facial expressions in adulthood. Despite these differences, our findings and those of these prior studies suggest that normative variability in caregiving and family functioning is associated with alterations in threat-related brain function similar to those that have been reported for caregiving extremes, including trauma, abuse, and neglect.

Because exaggerated threat-related amygdala reactivity has been linked to a range of clinical outcomes, particularly following stressful life events (Swartz, Williamson, et al., 2015), next steps could include probing the relationship between family functioning, clinical outcomes following stressors, and amygdala function. Although we found no main effect of familial affective responsiveness on concurrent symptoms of depression and anxiety in the present study, it is possible that the association with baseline amygdala reactivity may map onto future symptoms of stress-related anxiety and depression (Admon et al., 2013; Mattson et al.,

2016; McLaughlin et al., 2014; Swartz, Williamson, et al., 2015). Our initial evidence linking normative variability in caregiving with threat-related brain function can now serve as a starting point for pursuing longitudinal data analyses to investigate potential trajectories of amygdala reactivity and family functioning over time. Future research can further explore this question using whole-brain analyses, as well as examining functional connectivity between the amygdala and prefrontal cortex during the explicit regulation of negative emotions to gauge the impact of family functioning on top-down control processes.

Our study, of course, is not without limitations that can be addressed in future research. First, we relied on self-reported measures of caregiving using a single instrument. Similarly, we assessed symptoms of anxiety and depression using self-report inventories. Thus, our findings may be subject to reporting bias and are likely more representative of the perception of events rather than of objective events. Second, hemispheric lateralization of significant associations between caregiving and amygdala reactivity was not hypothesized and therefore awaits replication before further useful speculation. Third, we did not have an explicit measure of parental overprotection. Thus, our speculation regarding possible links between affective responsiveness and this aspect of family functioning previously linked with sensitization and maladaptive responses to stress outcomes requires explicit testing. Fourth, we evaluated amygdala reactivity to only threat-related facial expressions. It is possible that normative variability in affective responsiveness may also map onto individual differences in amygdala reactivity to reward-related social signals, namely happy facial expressions, or emotionally neutral facial expressions, and that these associations may reflect additional pathways of risk or resilience to psychopathology. Last, the TAOS is not a population-representative sample but rather selected based on familial risk for depression. Future studies in diverse samples are necessary to evaluate the extent to which our current findings are present more broadly and, consequently, useful for informing risk for psychopathology more generally.

Despite these limitations, our findings suggest that better familial affective responsiveness is associated with a paradoxical increase in amygdala reactivity to explicit signals of interpersonal threat (i.e., angry facial expressions), especially in adolescents reporting lower recent stressful life events. Better familial affective responsiveness may exacerbate neural reactivity to interpersonal threat—particularly when coupled with fewer external stressors—by limiting opportunities for the expression and subsequent self-regulation of negative emotions. Although our findings extend a rich literature on the effects of extremes in caregiving on emotional brain function to more subtle, normative variability, they also indicate that a more careful consideration of putatively protective factors in the expression of risk-related brain function may be necessary in the absence of caregiving extremes.

References

- Adams, R. B., Jr., Gordon, H. L., Baird, A. A., Ambady, N., & Kleck, R. E. (2003). Effects of gaze on amygdala sensitivity to anger and fear faces. *Science*, *300*, 1536. <http://dx.doi.org/10.1126/science.1082244>
- Admon, R., Lubin, G., Rosenblatt, J. D., Stern, O., Kahn, I., Assaf, M., & Hendler, T. (2013). Imbalanced neural responsivity to risk and reward indicates stress vulnerability in humans. *Cerebral Cortex*, *23*, 28–35. <http://dx.doi.org/10.1093/cercor/bhr369>
- Admon, R., Lubin, G., Stern, O., Rosenberg, K., Sela, L., Ben-Ami, H., & Hendler, T. (2009). Human vulnerability to stress depends on amygdala's predisposition and hippocampal plasticity. *Proceedings of the National Academy of Sciences of the United States of America*, *106*, 14120–14125. <http://dx.doi.org/10.1073/pnas.0903183106>
- Andreotti, C., Garrard, P., Venkatraman, S. L., & Compas, B. E. (2015). Stress-related changes in attentional bias to social threat in young adults: Psychobiological associations with the early family environment. *Cognitive Therapy and Research*, *39*, 332–342. <http://dx.doi.org/10.1007/s10608-014-9659-z>
- Angold, A., Costello, E. J., Messer, S. C., Pickles, A., Winder, F., & Silver, D. (1995). Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *International Journal of Methods in Psychiatric Research*, *5*, 237–249.
- Belsky, J., & de Haan, M. (2011). Annual research review: Parenting and children's brain development: The end of the beginning. *Journal of Child Psychology and Psychiatry*, *52*, 409–428. <http://dx.doi.org/10.1111/j.1469-7610.2010.02281.x>
- Bernstein, D. P., Ahluvalia, T., Pogge, D., & Handelsman, L. (1997). Validity of the Childhood Trauma Questionnaire in an adolescent psychiatric population. *Journal of the American Academy of Child & Adolescent Psychiatry*, *36*, 340–348. <http://dx.doi.org/10.1097/00004583-199703000-00012>
- Bernstein, D. P., Stein, J. A., Newcomb, M. D., Walker, E., Pogge, D., Ahluvalia, T., . . . Zule, W. (2003). Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse & Neglect*, *27*, 169–190. [http://dx.doi.org/10.1016/S0145-2134\(02\)00541-0](http://dx.doi.org/10.1016/S0145-2134(02)00541-0)
- Birmaher, B., Brent, D. A., Chiappetta, L., Bridge, J., Monga, S., & Baugher, M. (1999). Psychometric properties of the Screen for Child Anxiety Related Emotional Disorders (SCARED): A replication study. *Journal of the American Academy of Child & Adolescent Psychiatry*, *38*, 1230–1236. <http://dx.doi.org/10.1097/00004583-199910000-00011>
- Birmaher, B., Dahl, R. E., Williamson, D. E., Perel, J. M., Brent, D. A., Axelson, D. A., . . . Ryan, N. D. (2000). Growth hormone secretion in children and adolescents at high risk for major depressive disorder. *Archives of General Psychiatry*, *57*, 867. <http://dx.doi.org/10.1001/archpsyc.57.9.867>
- Bogdan, R., Williamson, D. E., & Hariri, A. R. (2012). Mineralocorticoid receptor Iso/Val (rs5522) genotype moderates the association between previous childhood emotional neglect and amygdala reactivity. *American Journal of Psychiatry*, *169*, 515–522. <http://dx.doi.org/10.1176/appi.ajp.2011.11060855>
- Bouton, M. E. (2007). *Learning and behavior: A contemporary synthesis*. Sunderland, MA: Sinauer.
- Bowlby, J. (1978). Attachment theory and its therapeutic implications. *Adolescent Psychiatry*, *6*, 5–33.
- Bowlby, J., & Zeanah, C. H. (1988). A secure base: Parent-child attachment and healthy human development. *Journal of Nervous and Mental Disease*, *178*(1), 62.
- Caldji, C., Diorio, J., & Meaney, M. J. (2003). Variations in maternal care alter GABA_A receptor subunit expression in brain regions associated with fear. *Neuropsychopharmacology*, *28*, 1950–1959. <http://dx.doi.org/10.1038/sj.npp.1300237>
- Callaghan, B. L., & Tottenham, N. (2016). The neuro-environmental loop of plasticity: A cross-species analysis of parental effects on emotion circuitry development following typical and adverse caregiving. *Neuropsychopharmacology*, *41*, 163–176. <http://dx.doi.org/10.1038/npp.2015.204>
- Canetti, L., Bachar, E., Galili-Weisstub, E., De-Nour, A. K., & Shalev, A. Y. (1997). Parental bonding and mental health in adolescence. *Adolescence*, *32*, 381–394.
- Cicchetti, D., & Curtis, W. J. (2015). The developing brain and neural plasticity: Implications for normality, psychopathology, and resilience.

- In D. Cicchetti & D. J. Cohen (Eds.), *Developmental psychopathology* (pp. 1–64). <http://dx.doi.org/10.1002/9780470939390.ch1>
- Ekman, P., & Friesen, W. V. (1976). *Pictures of Facial Affect* [Image database]. Palo Alto, CA: Consulting Psychologists Press.
- Epstein, N. B., Baldwin, L. M., & Bishop, D. S. (1983). The McMaster Family Assessment Device. *Journal of Marital and Family Therapy*, 9, 171–180. <http://dx.doi.org/10.1111/j.1752-0606.1983.tb01497.x>
- Etkin, A., & Wager, T. D. (2007). Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*, 164, 1476–1488. <http://dx.doi.org/10.1176/appi.ajp.2007.07030504>
- Fanselow, M. S., & Tighe, T. J. (1988). Contextual conditioning with massed versus distributed unconditional stimuli in the absence of explicit conditional stimuli. *Journal of Experimental Psychology: Animal Behavior Processes*, 14, 187–199. <http://dx.doi.org/10.1037/0097-7403.14.2.187>
- Francis, D. D., & Meaney, M. J. (1999). Maternal care and the development of stress responses. *Current Opinion in Neurobiology*, 9, 128–134. [http://dx.doi.org/10.1016/S0959-4388\(99\)80016-6](http://dx.doi.org/10.1016/S0959-4388(99)80016-6)
- Gee, D. G., Gabard-Durnam, L. J., Flannery, J., Goff, B., Humphreys, K. L., Telzer, E. H., . . . Tottenham, N. (2013). Early developmental emergence of human amygdala-prefrontal connectivity after maternal deprivation. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 15638–15643. <http://dx.doi.org/10.1073/pnas.1307893110>
- Goodwin, R. D., Fergusson, D. M., & Horwood, L. J. (2004). Early anxious/withdrawn behaviours predict later internalising disorders. *Journal of Child Psychology and Psychiatry*, 45, 874–883. <http://dx.doi.org/10.1111/j.1469-7610.2004.00279.x>
- Green, M., & Solnit, A. J. (1964). Reactions to the threatened loss of a child: A vulnerable child syndrome. *Pediatrics*, 34, 58–66.
- Herringa, R. J., Birn, R. M., Ruttle, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., & Essex, M. J. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 19119–19124. <http://dx.doi.org/10.1073/pnas.1310766110>
- Kabacoff, R. I., Miller, I. W., Bishop, D. S., Epstein, N. B., & Keitner, G. I. (1990). A psychometric study of the McMaster Family Assessment Device in psychiatric, medical, and nonclinical samples. *Journal of Family Psychology*, 3, 431–439. <http://dx.doi.org/10.1037/h0080547>
- Kaufman, J., Birmaher, B., Brent, D., Rao, U., Flynn, C., Moreci, P., . . . Ryan, N. (1997). Schedule for Affective Disorders and Schizophrenia for School-Age Children—Present and Lifetime version (K-SADS-PL): Initial reliability and validity data. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36, 980–988. <http://dx.doi.org/10.1097/00004583-199707000-00021>
- Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 62, 593–602. <http://dx.doi.org/10.1001/archpsyc.62.6.593>
- Malter Cohen, M., Jing, D., Yang, R. R., Tottenham, N., Lee, F. S., & Casey, B. J. (2013). Early-life stress has persistent effects on amygdala function and development in mice and humans. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 18274–18278. <http://dx.doi.org/10.1073/pnas.1310163110>
- Mattson, W. I., Hyde, L. W., Shaw, D. S., Forbes, E. E., & Monk, C. S. (2016). Clinical neuroprediction: Amygdala reactivity predicts depressive symptoms 2 years later. *Social Cognitive and Affective Neuroscience*, 11, 892–898. <http://dx.doi.org/10.1093/scan/nsw018>
- McLaughlin, K. A., Busso, D. S., Duys, A., Green, J. G., Alves, S., Way, M., & Sheridan, M. A. (2014). Amygdala response to negative stimuli predicts PTSD symptom onset following a terrorist attack. *Depression and Anxiety*, 31, 834–842. <http://dx.doi.org/10.1002/da.22284>
- McLaughlin, K. A., & King, K. (2015). Developmental trajectories of anxiety and depression in early adolescence. *Journal of Abnormal Child Psychology*, 43, 311–323. <http://dx.doi.org/10.1007/s10802-014-9898-1>
- McLaughlin, K. A., Peverill, M., Gold, A. L., Alves, S., & Sheridan, M. A. (2015). Child maltreatment and neural systems underlying emotion regulation. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54, 753–762. <http://dx.doi.org/10.1016/j.jaac.2015.06.010>
- Messer, S. C., Angold, A., Costello, E. J., Loeber, R., Van Kammen, W., & Stouthamer-Loeber, M. (1995). The development of a short questionnaire for use in epidemiological studies of depression in children and adolescents: Factor composition and structure across development. *International Journal of Methods in Psychiatric Research*, 5, 251–262. Retrieved from <http://devepi.duhs.duke.edu/messermfarticle.pdf>
- Overbeek, G., ten Have, M., Vollebergh, W., & de Graaf, R. (2007). Parental lack of care and overprotection: Longitudinal associations with DSM-III-R disorders. *Social Psychiatry and Psychiatric Epidemiology*, 42, 87–93. <http://dx.doi.org/10.1007/s00127-006-0115-6>
- Parker, G. (1983). *Parental overprotection: A risk factor in psychosocial development*. New York, NY: Grune & Stratton.
- Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, 9, 947–957. <http://dx.doi.org/10.1038/nrn2513>
- Pechtel, P., Lyons-Ruth, K., Anderson, C. M., & Teicher, M. H. (2014). Sensitive periods of amygdala development: The role of maltreatment in preadolescence. *NeuroImage*, 97, 236–244. <http://dx.doi.org/10.1016/j.neuroimage.2014.04.025>
- Penny, W. D., Friston, K. J., Ashburner, J. T., Kiebel, S. J., & Nichols, T. E. (Eds.). (2011). *Statistical parametric mapping: The analysis of functional brain images*. New York, NY: Elsevier.
- Romund, L., Raufelder, D., Flemming, E., Lorenz, R. C., Pelz, P., Gleich, T., . . . Beck, A. (2016). Maternal parenting behavior and emotion processing in adolescents—An fMRI study. *Biological Psychology*, 120, 120–125. <http://dx.doi.org/10.1016/j.biopsycho.2016.09.003>
- Saxbe, D., Del Piero, L. B., Immordino-Yang, M. H., Kaplan, J. T., & Margolin, G. (2016). Neural mediators of the intergenerational transmission of family aggression. *Development and Psychopathology*, 28, 595–606. <http://dx.doi.org/10.1017/S0954579415000528>
- Scher, C. D., Stein, M. B., Asmundson, G. J. G., McCreary, D. R., & Forde, D. R. (2001). The Childhood Trauma Questionnaire in a community sample: Psychometric properties and normative data. *Journal of Traumatic Stress*, 14, 843–857. <http://dx.doi.org/10.1023/A:1013058625719>
- Schlund, M. W., & Cataldo, M. F. (2010). Amygdala involvement in human avoidance, escape and approach behavior. *NeuroImage*, 53, 769–776. <http://dx.doi.org/10.1016/j.neuroimage.2010.06.058>
- Sheridan, M. A., Fox, N. A., Zeanah, C. H., McLaughlin, K. A., & Nelson, C. A., III. (2012). Variation in neural development as a result of exposure to institutionalization early in childhood. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 12927–12932. <http://dx.doi.org/10.1073/pnas.1200041109>
- Stein, D., Williamson, D. E., Birmaher, B., Brent, D. A., Kaufman, J., Dahl, R. E., . . . Ryan, N. D. (2000). Parent-child bonding and family functioning in depressed children and children at high risk and low risk for future depression. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39, 1387–1395. <http://dx.doi.org/10.1097/00004583-200011000-00013>
- Susman, E. J. (2006). Psychobiology of persistent antisocial behavior: Stress, early vulnerabilities and the attenuation hypothesis. *Neuroscience and Biobehavioral Reviews*, 30, 376–389. <http://dx.doi.org/10.1016/j.neubiorev.2005.08.002>

- Swartz, J. R., Knodt, A. R., Radtke, S. R., & Hariri, A. R. (2015). A neural biomarker of psychological vulnerability to future life stress. *Neuron*, *85*, 505–511. <http://dx.doi.org/10.1016/j.neuron.2014.12.055>
- Swartz, J. R., Waller, R., Bogdan, R., Knodt, A. R., Sabhlok, A., Hyde, L. W., & Hariri, A. R. (2017). A common polymorphism in a Williams syndrome gene predicts amygdala reactivity and extraversion in healthy adults. *Biological Psychiatry*, *81*, 203–210. <http://dx.doi.org/10.1016/j.biopsych.2015.12.007>
- Swartz, J. R., Williamson, D. E., & Hariri, A. R. (2015). Developmental change in amygdala reactivity during adolescence: Effects of family history of depression and stressful life events. *American Journal of Psychiatry*, *172*, 276–283. <http://dx.doi.org/10.1176/appi.ajp.2014.14020195>
- Tan, P. Z., Lee, K. H., Dahl, R. E., Nelson, E. E., Stroud, L. J., Siegle, G. J., . . . Silk, J. S. (2014). Associations between maternal negative affect and adolescent's neural response to peer evaluation. *Developmental Cognitive Neuroscience*, *8*, 28–39. <http://dx.doi.org/10.1016/j.dcn.2014.01.006>
- Taylor, S. E., Eisenberger, N. I., Saxbe, D., Lehman, B. J., & Lieberman, M. D. (2006). Neural responses to emotional stimuli are associated with childhood family stress. *Biological Psychiatry*, *60*, 296–301. <http://dx.doi.org/10.1016/j.biopsych.2005.09.027>
- Thijssen, S., Muetzel, R. L., Bakermans-Kranenburg, M. J., Jaddoe, V. W. V., Tiemeier, H., Verhulst, F. C., . . . van IJzendoorn, M. H. (2017). Insensitive parenting may accelerate the development of the amygdala-medial prefrontal cortex circuit. *Development and Psychopathology*, *29*, 505–518. <http://dx.doi.org/10.1017/S0954579417000141>
- Thomasgard, M., & Metz, W. P. (1993). Parental overprotection revisited. *Child Psychiatry and Human Development*, *24*, 67–80. <http://dx.doi.org/10.1007/BF02367260>
- Tottenham, N. (2012). Human amygdala development in the absence of species-expected caregiving. *Developmental Psychobiology*, *54*, 598–611. <http://dx.doi.org/10.1002/dev.20531>
- Tottenham, N., Hare, T. A., Quinn, B. T., McCarry, T. W., Nurse, M., Gilhooly, T., . . . Casey, B. J. (2010). Prolonged institutional rearing is associated with atypically large amygdala volume and difficulties in emotion regulation. *Developmental Science*, *13*, 46–61. <http://dx.doi.org/10.1111/j.1467-7687.2009.00852.x>
- Tottenham, N., Shapiro, M., Telzer, E. H., & Humphreys, K. L. (2012). Amygdala response to mother. *Developmental Science*, *15*, 307–319. <http://dx.doi.org/10.1111/j.1467-7687.2011.01128.x>
- Whalen, P. J., & Phelps, E. A. (Eds.). (2009). *The human amygdala*. New York, NY: Guilford Press.
- White, M. G., Bogdan, R., Fisher, P. M., Muñoz, K. E., Williamson, D. E., & Hariri, A. R. (2012). FKBP5 and emotional neglect interact to predict individual differences in amygdala reactivity. *Genes, Brain and Behavior*, *11*, 869–878. <http://dx.doi.org/10.1111/j.1601-183X.2012.00837.x>
- Whittle, S., Yap, M. B. H., Yücel, M., Sheeber, L., Simmons, J. G., Pantelis, C., & Allen, N. B. (2009). Maternal responses to adolescent positive affect are associated with adolescents' reward neuroanatomy. *Social Cognitive and Affective Neuroscience*, *4*, 247–256. <http://dx.doi.org/10.1093/scan/nsp012>
- Williamson, D. E., Birmaher, B., Axelson, D. A., Ryan, N. D., & Dahl, R. E. (2004). First episode of depression in children at low and high familial risk for depression. *Journal of the American Academy of Child & Adolescent Psychiatry*, *43*, 291–297. <http://dx.doi.org/10.1097/00004583-200403000-00010>
- Williamson, D. E., Birmaher, B., Ryan, N. D., Shiffrin, T. P., Lusk, J. A., Protopapa, J., . . . Brent, D. A. (2003). The stressful life events schedule for children and adolescents: Development and validation. *Psychiatry Research*, *119*, 225–241. [http://dx.doi.org/10.1016/S0165-1781\(03\)00134-3](http://dx.doi.org/10.1016/S0165-1781(03)00134-3)

Received October 5, 2017

Revision received April 12, 2018

Accepted April 24, 2018 ■